

Rabbit polyclonal antibody to Myristoylated alanine rich C kinase substrate (MARCKS): Whole serum

Catalogue No.:	R-1380-50
Description:	Myristoylated alanine-rich C-kinase substrate (MARCKS) is the most prominent cellular substrate for protein kinase C. This protein binds calmodulin, actin, and synapsin. MARCKS is a filamentous (F) actin cross-linking protein. Ref: SWISSPROT.
Batch No.:	See product label
Unit size:	50 uL
Antigen:	Recombinant full length human MARCKS expressed and purified from E. coli
Antibody Type:	Antiserum
Other Names:	Myristoylated alanine-rich C-kinase substrate; MARCKS; Protein kinase C substrate 80 kDa protein; Marcks; Macs;
Accession:	P30009 MARCS_RAT;
Produced in:	Rabbit
Applications:	Western Blotting (WB) and Immunocytochemistry (IC). A dilution of 1:1,000 - 1:2,000 is recommended for WB. A dilution of 1:500-1,000 is recommended for IC. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The specificity of this antibody has been confirmed by WB.
Species Against:	Human
Antibody Against:	Myristoylated alanine rich C kinase substrate (MARCKS)
Cross-reactivity:	Does not react with rodent protein.
Form:	Lyophilised
Reconstitution:	Reconstitute in 50 uL sterile distilled water. Centrifuge to remove any insoluble material.
Storage:	After reconstitution of lyophilised antibody, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
Expiry Date:	12 months after purchase




Image of mixed neuron/glia cultures stained with Rabbit polyclonal antibody to Myristoylated alanine rich C kinase substrate (MARCKS) R-1380-50 (red) and Chicken polyclonal antibody to MAP2 C-1382-50. Note that the MARCKS antibody stains vesicular structures both in the glial cells and in the dendrites of the neurons, which are strongly stained with the MAP2 antibody.

FOR RESEARCH USE ONLY