



Mouse monoclonal antibody to Visinin-like protein 1


Catalogue No.:	M-1657-100
Description:	Visinin (sometimes known as hippocalcin-like protein 3, HLP3, HPCAL3, HUVISL1, VLP-1, VILIP and VILIP-1) was originally isolated biochemically from chicken retina as a major protein of about 24kDa on SDS-PAGE (1). Following cloning and sequencing of visinin, several visinin like proteins were discovered by homology screening (2, 3). One of these, Visinin-like protein 1 is a small Calcium binding protein which is very abundant in the nervous system and is found only in neurons, though different neurons have different levels of expression (4, 5). It is particularly concentrated in cerebellar Purkinje cells, and tends to be most abundant in perikarya and dendrites. The protein belongs to the large superfamily of calmodulin and parvalbumin type proteins which function by binding Calcium ions. Calcium binding alters the conformation of these proteins and allow them to interact with other binding partners, the properties of which they may alter. Visinin-like protein 1 has four "EF hand" domains, which are negatively charged helix-turn-helix peptides which are responsible for Calcium binding. Visinin-like protein 1 is 191 amino acids in size and has a molecular weight on SDS-PAGE of 22kDa. The protein has recently been suggested to be a useful biomarker of Alzheimer's disease and traumatic brain injury (6, 7, 8).
Unit size:	100 ug
Antigen:	Recombinant Visinin-like protein 1 expressed and purified from E. coli.
Antibody Type:	Monoclonal
Isotype:	IgG1
Other Names:	Hippocalcin-like protein 3, HLP3, HPCAL3, HUVISL1, VLP-1, VILIP and VILIP-1
Produced in:	Mouse
Applications:	Western Blotting (WB) and Immunocytochemistry (IC). A dilution of 1:1,000 - 1:2,000 is recommended for WB. A dilution of 1:500-1:1,000 is recommended for IC. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The antibody reacts with a 22 kDa band by Western blot on bovine cerebellum lysate. It has also been used successfully for immunocytochemistry.
Species Against:	Human, bovine, rat and mouse. It is expected that it will work on other mammal tissues.
Antibody Against:	Visinin-like protein 1
Form:	Lyophilised from PBS. Contains 5% trehalose.
Appearance:	White powder
Reconstitution:	Reconstitute in sterile distilled water. Centrifuge to remove any insoluble material.
Storage:	After reconstitution of lyophilised antibody, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
Expiry Date:	12 months after purchase
General References:	1. Hatakenaka S, Kuo CH, Miki N. Analysis of a distinctive protein in chick retina during development. Brain Res. 312:155-63 (1983).

FOR RESEARCH USE ONLY



Mouse monoclonal antibody to Visinin-like protein 1

2. Kuno T, Kajimoto Y, Hashimoto T, Mukai H, Shirai Y, Saheki S, Tanaka C. cDNA cloning of a neural visinin-like Ca(2+)-binding protein. *Biochem Biophys Res Commun.* 184:1219-25 (1992).
3. Polymeropoulos MH, Ide S, Soares MB, Lennon GG. Sequence characterization and genetic mapping of the human VSNL1 gene, a homologue of the rat visinin-like peptide RNVP1. *Genomics.* 29:273-5 (1995).
4. Bernstein HG, Baumann B, Danos P, Diekmann S, Bogerts B, Gundelfinger ED, Braunewell KH. Regional and cellular distribution of neural visinin-like protein immunoreactivities (VILIP-1 and VILIP-3) in human brain. *J Neurocytol.* 28:655-62 (1999).
5. Paterlini M, Revilla V, Grant AL, Wisden W. Expression of the neuronal calcium sensor protein family in the rat brain. *Neuroscience.* 99:205-16 (2000)
6. Laterza OF, Modur VR, Crimmins DL, Olander JV, Landt Y, Lee JM, Ladenson JH. Identification of novel brain biomarkers. *Clin. Chem.* 9:1713-21 (2006)
7. Lee JM, Blennow K, Andreasen N, Laterza O, Modur V, Olander J, Gao F, Ohlendorf M, Ladenson JH. The brain injury biomarker VLP-1 is increased in the cerebrospinal fluid of Alzheimer disease patients. *Clin. Chem.* 10:1617-23 (2008).
8. Tarawneh R, D'Angelo G, Macy E, Xiong C, Carter D, Cairns NJ, Fagan AM, Head D, Mintun MA, Ladenson JH, Lee JM, Morris JC, Holtzman DM. Visinin-like protein-1: diagnostic and prognostic biomarker in Alzheimer disease. *Ann Neurol.* 70:274-85 (2011) doi: 10.1002/ana.22448.



```
/* Font Definitions */
@font-face
{font-family:"Times New Roman";
panose-1:0 2 2 6 3 5 4 5 2 3;
mso-font-charset:0;
mso-generic-font-family:auto;
mso-font-pitch:variable;
mso-font-signature:50331648 0 0 0 1 0;}
@font-face
{font-family:Arial;
panose-1:0 2 11 6 4 2 2 2 2 2;
mso-font-charset:0;
mso-generic-font-family:auto;
mso-font-pitch:variable;
mso-font-signature:50331648 0 0 0 1 0;}
@font-face
{font-family:"Lucida Grande";
mso-font-charset:0;
```

FOR RESEARCH USE ONLY



Mouse monoclonal antibody to Visinin-like protein 1

```
mso-generic-font-family:auto;
mso-font-pitch:variable;
mso-font-signature:50331648 0 0 0 1 0;}
@font-face
{font-family:Tahoma;
panose-1:0 2 11 6 4 3 5 4 4 2;
mso-font-charset:0;
mso-generic-font-family:auto;
mso-font-pitch:variable;
mso-font-signature:50331648 0 0 0 1 0;}
/* Style Definitions */
p.MsoNormal, li.MsoNormal, div.MsoNormal
{mso-style-parent:"";
margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
font-size:10.0pt;
font-family:"Times New Roman";}
h1
{mso-style-next:Normal;
margin-top:0in;
margin-right:0in;
margin-bottom:0in;
margin-left:-.9pt;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
page-break-after:avoid;
mso-outline-level:1;
font-size:12.0pt;
font-family:Arial;
mso-font-kerning:0pt;
font-weight:normal;}
h2
{mso-style-next:Normal;
margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
page-break-after:avoid;
mso-outline-level:2;
font-size:12.0pt;
font-family:Arial;}
h3
{mso-style-next:Normal;
margin:0in;
margin-bottom:.0001pt;
text-align:center;
mso-pagination:widow-orphan;
page-break-after:avoid;
mso-outline-level:3;
font-size:12.0pt;
font-family:"Lucida Grande";
color:black;}
```

FOR RESEARCH USE ONLY



Mouse monoclonal antibody to Visinin-like protein 1

```
p.MsoHeader, li.MsoHeader, div.MsoHeader
{margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
tab-stops:center 3.0in right 6.0in;
font-size:10.0pt;
font-family:"Times New Roman";}
p.MsoFooter, li.MsoFooter, div.MsoFooter
{margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
tab-stops:center 3.0in right 6.0in;
font-size:10.0pt;
font-family:"Times New Roman";}
p.MsoTitle, li.MsoTitle, div.MsoTitle
{margin:0in;
margin-bottom:.0001pt;
text-align:center;
mso-pagination:widow-orphan;
font-size:14.0pt;
font-family:"Times New Roman";}
p.MsoBodyText, li.MsoBodyText, div.MsoBodyText
{margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
font-size:11.0pt;
font-family:Arial;}
p.MsoBodyTextIndent, li.MsoBodyTextIndent, div.MsoBodyTextIndent
{margin-top:0in;
margin-right:0in;
margin-bottom:0in;
margin-left:.5in;
margin-bottom:.0001pt;
text-indent:-.5in;
line-height:12.0pt;
mso-line-height-rule:exactly;
mso-pagination:widow-orphan;
tab-stops:135.9pt;
font-size:9.0pt;
font-family:Arial;}
p.MsoSubtitle, li.MsoSubtitle, div.MsoSubtitle
{margin:0in;
margin-bottom:.0001pt;
text-align:center;
mso-pagination:widow-orphan;
font-size:14.0pt;
font-family:Arial;
font-weight:bold;}
p.MsoBodyText2, li.MsoBodyText2, div.MsoBodyText2
{margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
```

FOR RESEARCH USE ONLY



Mouse monoclonal antibody to Visinin-like protein 1

```
font-size:8.0pt;
font-family:Arial;
color:black;}
p.MsoBodyText3, li.MsoBodyText3, div.MsoBodyText3
{margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
font-size:11.0pt;
font-family:Arial;
color:black;}
p.MsoBodyTextIndent2, li.MsoBodyTextIndent2, div.MsoBodyTextIndent2
{margin-top:0in;
margin-right:0in;
margin-bottom:0in;
margin-left:9.0pt;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
font-size:9.0pt;
font-family:Arial;
color:black;}
p.MsoBodyTextIndent3, li.MsoBodyTextIndent3, div.MsoBodyTextIndent3
{margin-top:0in;
margin-right:0in;
margin-bottom:0in;
margin-left:9.0pt;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
font-size:11.0pt;
font-family:Arial;
color:black;}
p.BalloonText, li.BalloonText, div.BalloonText
{mso-style-name:"Balloon Text";
margin:0in;
margin-bottom:.0001pt;
mso-pagination:widow-orphan;
font-size:8.0pt;
font-family:Tahoma;}
@page Section1
{size:8.5in 11.0in;
margin:1.0in 1.25in 1.0in 1.25in;
mso-header-margin:.5in;
mso-footer-margin:.5in;
mso-paper-source:0;}
div.Section1
{page:Section1;}
/* List Definitions */
@list l0
{mso-list-id:1970355742;
mso-list-type:hybrid;
mso-list-template-ids:1602914926 1555098738 1639433 1770505
984073 1639433 1770505 984073 1639433 1770505;}
@list l0:level1
```

FOR RESEARCH USE ONLY



Mouse monoclonal antibody to Visinin-like protein 1

```
{mso-level-tab-stop:27.75pt;  
mso-level-number-position:left;  
margin-left:27.75pt;  
text-indent:-.25in;}
```

ol

```
{margin-bottom:0in;}
```

ul

```
{margin-bottom:0in;}
```

-->

color:black;mso-ansi-language:EN-US">Western blot of bovine cerebellum homogenate stained with M-1657-100.

biosensis

FOR RESEARCH USE ONLY