



Mouse monoclonal antibody to High-mobility group protein box 1 (HMGB1): IgG

Catalogue No.:	M-1702-100
Description:	High-mobility group proteins were named originally since they are abundant relatively low molecular weight proteins which run quickly on SDS-PAGE gels. High-mobility group protein box 1 (HMGB1, Amphoterin) is one of these. The "bx" in the name refers to the so-called high mobility group (HMG) box, a compact domain involved in DNA binding and protein-protein interactions. the HMGB1 molecule has two of these HMG domains. The protein is also called amphoterin, this name being derived from the presence of two highly charged regions in the molecule, a relatively neutrally charged N-terminus and a very negatively charged C-terminus. In fact the molecule is very unusually charged throughout, the human sequence consisting of 16.7% Glutamic acid, 9.3% Aspartic acid, 20% lysine and 9.3% Arginine. HMGB1 can bind Toll like receptor 4 (TLR4) and the Receptor for Advanced Glycation End products (RAGE). TLRs are components of the innate immune system, first recognized as a family of receptors which recognize "Pathogen Associated Molecular Pattern molecules (PAMPs). PAMPs are common components of bacteria and when TLRs bind these a strong inflammatory response is activated. More recently it has been recognized that TLRs can also be activated by Damage Associated Molecular Pattern molecules (DAMPs), which are endogenous substances released from damaged and diseased cells which also bind to TLR family receptors and also activate inflammation. HMGB1 is such a DAMP, binding to TLR4, and much evidence suggests that HMGB1 is a strong activator of inflammation. Interestingly, HMGB1 is released by necrotic cells but not by apoptotic cells (1).
Batch No.:	See product label
Unit size:	100 uL
Antigen:	HMGB1
Antibody Type:	Monoclonal
Isotype:	IgG2b
Produced in:	Mouse
Applications:	Western Blotting (WB), Immunocytochemistry (ICC) and Flow Cytometry. A dilution of 1:1,000 - 1:2,000 is recommended for WB. A dilution of 1:500 - 1:1,000 is recommended for ICC. For Flow Cytometry, use ~2 ug per 10 ⁶ cells. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The antibody reacts with a band at ~25 kDa by Western blot on HeLa cell extract. It has also been used successfully for immunocytochemistry showing strong nuclear staining.
Species Against:	Human, bovine, porcine, rat and mouse. It is expected that it will work on other mammal tissues.
Form:	Lyophilised purified culture supernatant with 5% trehalose and 0.5% sodium azide.
Appearance:	White powder
Reconstitution:	Reconstitute in sterile distilled water. Centrifuge to remove any insoluble material.


FOR RESEARCH USE ONLY

Mouse monoclonal antibody to High-mobility group protein box 1 (HMGB1): IgG


Storage: After reconstitution of lyophilised antibody, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.

Expiry Date: 12 months after purchase

References: 1. Scaffidi P, Misteli T, Bianchi ME. Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* 418:191-195 (2002).



B



A: HeLa cells stained with M-1702-100 (red), chicken polyclonal antibody to Vimentin (C-1409-50, green) and DNA (blue). The M-1702-100 antibody reveals strong nuclear staining which overlaps with the DNA stain. B: Blot of crude HeLa cell extract stained with M-1702-100. HMGB1 runs at an apparent molecular weight of 25 kDa. C: Analysis of HMGB1 expression in human euroblastoma SH-SY5Y cell line by Flow Cytometry. Fixing and Permeabilization of cells: Absolute methanol (10 minutes in ice) and 0.1% Tween-20 in PBS, Blocking: 1% BSA, Primary antibody: Mouse Monoclonal antibody to HMGB1 (cat # M-1702-100, 2 μ g per $\sim 10^6$ cells) for 30 minutes at room temperature, Secondary antibody: Goat anti-mouse PE labeled secondary antibody (1:100 fold dilution) with incubation for 20 minutes in dark at room temperature. Non-specific Control IgG, clone X63 (cat # M-1249-200) was used as negative control under same conditions (black dashed). Flow cytometry data and results were generated using Orflo MoxiflowTM instrument and protocols.

FOR RESEARCH USE ONLY