

Non-specific monoclonal IgG control antibody [X63], Saporin-conjugated

Catalogue No.: M-1792-50-IT

Description: THIS PRODUCT IS TEMPORARILY OUT OF STOCK. PLEASE REFER TO THE "REPLACED

> BY" FIELD BELOW TO LOCATE THE CURRENT BIOSENSIS PRODUCT TO MEET YOUR RESEARCH NEEDS. X63-saporin is an antibody conjugate comprising of the non-specific monoclonal IgG1 antibody X63, chemically conjugated via a reducible disulfide bridge to the ribosome-inactivating protein saporin, purified from saponaria officinalis. Antibody clone X63 has no known binding ability, and thus can be used as negative control antibody. In combination with saporin, X63-saporin is a useful negative control for targeting IgG1-saporin

conjugates such as MC192-saporin.

Related products: M-1755-50-IT, Mouse monoclonal antibody to rat p75NTR [MC192], Saporin-conjugated, 50

μg

M-1249-100, Non-specific control IgG [X63]: IgG1

PE-001-100, Purified Saporin

R-025-100, Rabbit antibody to Saporin: whole serum

Batch No.: See product label.

Unit size: 50 µg

Antigen: None. This control IgG has no known binding ability.

Isotype: lgG1 Clone: X63

Accession: RIP6_SAPOF

Molecular Weight: The average molecular weight of the X63-saporin conjugate is approximately 210 kDa.

Handling: Only qualified personnel should handle this product. X63 antibody has no known target

> receptor, however, standard laboratory protective measures should be followed, including wearing lab coat, gloves and safety glasses. Avoid contact with open wounds and practice good laboratory technique to avoid auto-exposure/auto-injection! Avoid contact of conjugate with reducing agents (eg., DTT, beta-mercaptoethanol) as this will destroy the antibody-saporin

conjugate.

Purity: > 90% by non-reducing SDS-PAGE

Biol. activity: None. Routinely tested for absence of killing of rat C6 cells in vitro. Note that the primary use of

X63-saporin is for in vivo applications.

Applications: Negative control for targeting IgG1-saporin conjugates., for instance MC192-saporin.

X63-saporin should be used at a concentration equal to that of the test antibody-saporin

conjugate.

Specificity: No staining has ever been identified with this immunoglobulin demonstrating its non-specific

value as a control.

Cross-reactivity: N/A

Conjugate: X63 antibody is chemically linked to saporin via a disulfide bridge.

FOR RESEARCH USE ONLY



Non-specific monoclonal IgG control antibody [X63], Saporin-conjugated

Form: Liquid. Provided as 1 mg/mL conjugate in PBS, pH 7.2-7.6, without preservative.

Storage: Product is shipped on ice packs and may arrive thawed. Upon receipt, pulse-centrifuge the vial

> to collect any liquid that may be entrapped in the lid. Immediately prepare aliquots and keep the undiluted stock at -80°C for long-term storage. Avoid repeated thaw-freezing. For short-term storage, keep at 2-8°C for up to 2 weeks. it is recommended to handle this product under

sterile conditions.

Expiry Date: 12 months from receipt when stored at -80°C (undiluted conjugate only).

