

Sheep antibody to rh NT3: IgG

Catalogue No.:	S-057-500
Description:	NT3 is a member of the neurotrophin family, that controls survival and differentiation of visceral and proprioceptive sensory neurons. NT3 is closely related to both NGF and BDNF. It may be involved in the maintenance of the adult nervous system, and may affect development of neurons in the embryo when it is expressed in human placenta. NT3-deficient mice generated by gene targeting display severe movement defects of the limbs. The mature peptide of this protein is identical in all mammals examined including human, pig, rat and mouse. SUBCELLULAR LOCATION: Secreted protein. TISSUE SPECIFICITY: Brain and peripheral tissues. SIMILARITY: Belongs to the NGF-beta family.
Batch No.:	See product label
Unit size:	500 ug
Antigen:	Recombinant human NT3
Other Names:	Neurotrophin-3; Neurotrophic factor; HDNF; Nerve growth factor 2; NGF-2; Ntf3; Ntf-3
Accession:	NT3_HUMAN
Produced in:	Sheep
Purity:	Protein G purified IgG
Applications:	IHC, ELISA, WB, inhibition of biological activity. A concentration of 1-10 ug/mL is recommended for IHC, ELISA, WB and inhibition of biological activity in vitro; 2-10 ug/mL (ED50) for in vivo use. This antiserum will neutralise NT3 but not other neurotrophins. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	A cross reactivity of less than 1% to mouse NGF, recombinant human BDNF and 5% to NT4/5 has been shown by 1-site ELISA.
Cross-reactivity:	This antiserum is known to react with rat, chicken and human NT3.
Form:	Lyophilised. Antibody is carrier-free. It contain no BSA or any other protein or sugar added. The material has been lyophilized from simple PBS buffer (10 mM phosphate, 150 mM NaCl, pH 7.2-7.6) without addition of sodium azide or other preservatives.
Reconstitution:	Reconstitute in 500 uL of sterile water. Centrifuge to remove any insoluble material.
Storage:	After reconstitution keep aliquots at -20C for a higher stability, and at 2-8C with an appropriate antibacterial agent. Glycerol (1:1) may be added for an additional stability. Avoid repetitive freeze/thaw cycles.
Expiry Date:	12 months after purchase
References:	1. Zouh et al. (1996) Mol. Neurobio. 13(3): 185-97 2. Rush et al. (1997) Clin. Exp. Pharmacol. 24: 549-55 3. Tafreshi et al. (1998) Neurosci. 83(2): 373-80

FOR RESEARCH USE ONLY