



Rabbit antibody to SUMO-1 (6-21): whole serum


Catalogue No.:	R-115-100
Description:	FUNCTION: Ubiquitin-like protein which binds to a wide range of target proteins. Does not seem to be involved in protein degradation and may function as an antagonist of ubiquitin in the degradation process. Plays a role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction. Involved in targeting RANGAP1 to the nuclear pore complex protein RANBP2. SUBUNIT: Covalently attached to a number of proteins such as PML, RANGAP1, HIPK2, SP100, p53, p73alpha, MDM2, JUN and DNMT3B. Also interacts with HIF1A, HIPK2, HIPK3, CHD3, PIAS1, EXOSC9, TDG, RAD51 and RAD52. SUBCELLULAR LOCATION: Nucleus; nuclear membrane. Nucleus; nucleoplasm; nuclear speckle. Cytoplasm. SIMILARITY: Belongs to the ubiquitin family. SMT3 subfamily. SIMILARITY: Contains 1 ubiquitin-like domain.
Batch No.:	See product label
Unit size:	100 uL
Antigen:	A synthetic peptide (AKPSTEDLGDKKEGEY) as part of human SUMO-1 peptide (aa: 6-21) conjugated to diphtheria toxoid has been used as the immunogen. This antigen is homologous with SUMO-1 of rat.
Other Names:	Small ubiquitin-related modifier 1; Ubiquitin-like protein SMT3C; SMT3 homolog 3; Ubiquitin-homology domain protein PIC1; Ubiquitin-like protein UBL1; GAP-modifying protein 1; GMP1; Sentrin; SUMO1; SMT3C; SMT3H3; UBL1
Accession:	SUMO1_HUMAN
Produced in:	Rabbit
Purity:	Whole serum
Applications:	IHC, WB. A dilution of 1:1000 to 1:2000 is recommended for immunohistochemistry and 1:2000 to 1:4000 for western blot. Cell lysate from Hela and NIH-3T3 cell lysates may be used as a positive control, and for IHC, lung carcinoma may be used. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	This antiserum recognises human SUMO-1 and not ubiquitin.
Cross-reactivity:	This antiserum is known to cross react with rat and human SUMO-1.
Form:	Lyophilised
Reconstitution:	Reconstitute in 100 uL of sterile water. Centrifuge to remove any insoluble material.
Storage:	After reconstitution keep aliquots at -20C for a higher stability, and at 2-8C with an appropriate antibacterial agent. Glycerol (1:1) may be added for an additional stability. Avoid repetitive freeze/thaw cycles.
Expiry Date:	12 months after purchase
References:	1. Yang, S.H. et al. Mol. Cell 13(4):611-617 (2004). 2. Ohshima, T. et al. J. Biol. Chem. 278(51):50833-50842 (2003). 3. Bailey, D. et al. J. Biol. Chem. 279(1):692-703 (2004). 4. Ling, Y. et al. Nucleic Acids Res. 32(2):598-610 (2004).

FOR RESEARCH USE ONLY



Rabbit antibody to SUMO-1 (6-21): whole serum

5. Pountney, D.L. et al. Exp. Neurol. 184(1):436-446 (2003).



Western blot detection of recombinant SUMO-1 protein using Rabbit antibody to SUMO-1 (6-21): ws (catalogue no. R-115-100) at a dilution of 1:4000. The protein sample was 1 μ l of E.Coli cell lysate containing approximately 50 ng of recombinant SUMO-1 protein. The antibody detects a band of 17 kDa.

biosensis

FOR RESEARCH USE ONLY