



## Rabbit antibody to the Tyrosine Kinase Receptor C (TrkC): whole serum

<b>Catalogue No.:</b>	R-151-100
<b>Description:</b>	TrkC is a member of the neurotrophic tyrosine receptor kinase family. TrkC is a membrane-bound receptor that upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. TrkC is the receptor for neurotrophin-3 (NT-3).? Signalling through TrkC leads to cell differentiation and may play a role in the development of proprioceptive neurons that sense body position. SUBUNIT: Exists in a dynamic equilibrium between monomeric (low affinity) and dimeric (high affinity) structures. SUBCELLULAR LOCATION: Membrane; single-pass type I membrane protein. ALTERNATIVE PRODUCTS: 4 named isoforms produced by alternative splicing. Additional isoforms seem to exist. Mutations in TrkC have been associated with medulloblastomas, secretory breast carcinomas and other cancers.
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 uL
<b>Antigen:</b>	Extracellular domain of glycosylated human TrkC protein produced in CHO cells was used as the immunogen.
<b>Other Names:</b>	Tropomyosin-related kinase receptor; NT-3 growth factor receptor; Neurotrophic tyrosine kinase receptor type 3; TrkC tyrosine kinase; GP145-TrkC; Trk-C; NTRK3; TRKC
<b>Accession:</b>	NTRK3_HUMAN
<b>Produced in:</b>	Rabbit
<b>Purity:</b>	Whole serum
<b>Applications:</b>	IHC. A dilution of 1:1000 to 1:3000 is recommended for this application. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	Specificity was demonstrated by immunohistochemistry. This antibody was used to stain cryostat sections of the rat peripheral sensory ganglia.
<b>Cross-reactivity:</b>	Reacts with human, rat and mouse TrkC. Other species have not yet been tested.
<b>Form:</b>	Lyophilised
<b>Reconstitution:</b>	Reconstitute in 100 uL of sterile water. Centrifuge to remove any insoluble material.
<b>Storage:</b>	After reconstitution keep aliquots at -20C for a higher stability, and at 2-8C with an appropriate antibacterial agent. Glycerol (1:1) may be added for an additional stability. Avoid repetitive freeze/thaw cycles.
<b>References:</b>	Zhang FX, et al. (2005) Brain Res. 1062 (1-2) pp. 92-100.

---

FOR RESEARCH USE ONLY