



## Rabbit polyclonal antibody to Catenin gamma (8-22): Affinity purified

<b>Catalogue No.:</b>	R-1027-100
<b>Description:</b>	Catenin gamma, also known as Junction Plakoglobin, is a common junctional plaque protein. It is a cytoplasmic protein with soluble and membrane-associated forms. The presence of Catenin gamma in both the desmosomes and in the intermediate junctions suggests it has a role in the structure and function of submembranous plaques.
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 ug
<b>Antigen:</b>	A synthetic peptide (EQPIKVTEWQQTYTY) corresponding to a region (8-22) from the N-terminus of human Catenin gamma.
<b>Other Names:</b>	Junction plakoglobin; Desmoplakin; JUP; CTNNG; DP3;
<b>Accession:</b>	P14923 PLAK_HUMAN;
<b>Produced in:</b>	Rabbit
<b>Purity:</b>	Affinity purified on antigen column
<b>Applications:</b>	Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 1.0 ug/mL is recommended for WB. Human Catenin gamma has a predicted length of 745 residues and MW of 82 kDa. A concentration of 1.0-2.0 ug/mL is recommended to detect Catenin gamma in formalin fixed and paraffin embedded tissues. Antigen retrieval is required. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
<b>Cross-reactivity:</b>	Human; mouse; rat;
<b>Form:</b>	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg NaN <sub>3</sub>
<b>Reconstitution:</b>	Reconstitute in 100 uL of sterile distilled water to achieve an antibody concentration of 1 mg/mL. Centrifuge to remove any insoluble material.
<b>Storage:</b>	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
<b>Expiry Date:</b>	12 months after purchase

---

FOR RESEARCH USE ONLY