

## Rabbit polyclonal antibody to human NTKL-binding protein 1 (270-285): Affinity purified

<b>Catalogue No.:</b>	R-1077-100
<b>Description:</b>	NTKL-binding protein 1 (GORAB) belongs to the golgin family of coiled-coil proteins located in the golgi apparatus. GORAB interacts with SCYL1, RCHY1 and RAB6A/RAB6. At least 4 isoforms are produced by alternative splicing.
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 ug
<b>Antigen:</b>	A synthetic peptide (KEQLTEHLCTIIQQNE) corresponding to a region (270-285) from human NTKL-binding protein 1 (GORAB). To enhance the immunological response, this peptide was coupled to carrier protein BSA.
<b>Other Names:</b>	RAB6-interacting golgin; N-terminal kinase-like-binding protein 1; NTKL-binding protein 1; NTKL-BP1; hNTKL-BP1; SCY1-like 1-binding protein 1; SCYL1-binding protein 1; SCYL1-BP1; GORAB; NTKLBP1; SCYL1BP1; FLJ11752; MGC51263; MGC70512;
<b>Accession:</b>	Q5T7V8 GORAB_HUMAN;
<b>Produced in:</b>	Rabbit
<b>Purity:</b>	Affinity purified on antigen column
<b>Applications:</b>	Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 0.5-1.0 ug/mL is recommended for WB. Human NTKL-binding protein 1 (isoform 1) has a predicted length of 394 residues and MW of 45 kDa. A concentration of 1.0-2.0 ug/mL is recommended to detect the protein in formalin fixed and paraffin embedded tissues. Antigen retrieval is required. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
<b>Cross-reactivity:</b>	Human; mouse; rat; rabbit;
<b>Form:</b>	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg NaN <sub>3</sub>
<b>Reconstitution:</b>	Reconstitute in 100 uL of sterile distilled water to achieve an antibody concentration of 1 mg/mL. Centrifuge to remove any insoluble material.
<b>Storage:</b>	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles
<b>Expiry Date:</b>	12 months after purchase

---

FOR RESEARCH USE ONLY