

Rabbit polyclonal antibody to human Matrix metalloproteinase-14 (561-577): Affinity purified

Catalogue No.:	R-1090-100
Description:	THIS PRODUCT IS TEMPORARILY OUT OF STOCK. PLEASE REFER TO THE "REPLACED BY" FIELD BELOW TO LOCATE THE CURRENT BIOSENSIS PRODUCT TO MEET YOUR RESEARCH NEEDS. The matrix metalloproteinases (MMPs) are a large family of zinc endopeptidases. All MMPs are synthesized as inactive proenzymes. The activation of these proenzymes is a critical step that leads to degradation of extracellular matrix components such as fibronectin and collagen type III. MMP14 is expressed in stromal cells of colon, breast, and head and neck and activates MMP2 (progelatinase A) by cleavage.
Batch No.:	See product label
Unit size:	100 ug
Antigen:	A synthetic peptide (FFRRHGTPRRLLYCQRS) corresponding to a region (561-577) from human Matrix metalloproteinase-14. To enhance the immunological response, this peptide was coupled to carrier protein BSA.
Other Names:	MMP-14; EC 3.4.24.80; Membrane-type matrix metalloproteinase 1; MT-MMP 1; MTMMP1; Membrane-type-1 matrix metalloproteinase; MT1-MMP; MT1MMP; MMP-X1; MMP14;
Accession:	P50281 MMP14_HUMAN;
Produced in:	Rabbit
Purity:	Affinity purified on antigen column
Applications:	Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 0.1-0.5 ug/mL is recommended for WB. Human MMP14 has a predicted length of 582 residues and MW of 66 kDa. A concentration of 0.5-1.0 ug/mL is recommended to detect the protein in formalin fixed and paraffin embedded tissues as well as formalin or acetone fixed frozen tissues. Heat mediated antigen retrieval is required. A concentration of 0.5-1.0 ug/mL is recommended for IHC-F. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
Cross-reactivity:	Human; rat; expected to react with mouse due to sequence homology
Form:	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃
Reconstitution:	Reconstitute in 100 uL of sterile distilled water to achieve an antibody concentration of 1 mg/mL. Centrifuge to remove any insoluble material.
Storage:	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles
Expiry Date:	12 months after purchase

FOR RESEARCH USE ONLY