



Rabbit polyclonal antibody to human p27 KIP 1 (180-194): Affinity purified

Catalogue No.:	R-1121-100
Description:	Cyclin-dependent kinase inhibitor p27 (p27 KIP 1) is cell cycle regulatory mitotic inhibitor of cdk activity and an important regulator of cell cycle progression.
Batch No.:	See product label
Unit size:	100 ug
Antigen:	A synthetic peptide (NAGSVEQTPKKPGLR) corresponding to a region (180-194) from human p27 KIP 1. To enhance the immunological response, this peptide was coupled to carrier protein BSA.
Other Names:	Cyclin-dependent kinase inhibitor 1B; Cyclin-dependent kinase inhibitor p27; p27Kip1; CDKN1B; KIP1;
Accession:	P46527 CDN1B_HUMAN;
Produced in:	Rabbit
Purity:	Affinity purified on antigen column
Applications:	Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 0.1-0.5 ug/mL is recommended for WB. Human p27 KIP 1 has a predicted length of 198 residues and MW of 22 kDa. A concentration of 0.5-1.0 ug/mL is recommended to detect the protein in formalin fixed and paraffin embedded tissues. Heat mediated antigen retrieval is required. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
Cross-reactivity:	Human; rat; predicted to react with mouse due to sequence homology;
Form:	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃
Reconstitution:	Reconstitute in 100 uL of sterile distilled water to achieve an antibody concentration of 1 mg/mL. Centrifuge to remove any insoluble material.
Storage:	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles
Expiry Date:	12 months after purchase

FOR RESEARCH USE ONLY