


## Rabbit polyclonal antibody to Hypoxia-inducible factor-1 alpha (775-790): Affinity purified

<b>Catalogue No.:</b>	R-1061-100
<b>Description:</b>	THIS PRODUCT HAS BEEN SUPERCEDED. PLEASE REFER TO THE "REPLACED BY" FIELD BELOW TO LOCATE THE CURRENT BIOSENSIS PRODUCT TO MEET YOUR RESEARCH NEEDS. Hypoxia-inducible factor-1 alpha (HIF-1 alpha) is a transcription factor involved in the induction of oxygen regulated genes in response to hypoxia. HIF-1 alpha binds to a specific DNA sequence within the hypoxia response element of target gene promoters. HIF-1 alpha is expressed in most tissue types and overexpressed in human cancers and their metastases. At least 2 isoforms of HIF-1 alpha are produced by alternative splicing.
<b>Replaced by:</b>	R-1722-100
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 µg
<b>Antigen:</b>	A synthetic peptide (PSDLACRLLGQSMDES) corresponding to a region (775-790) from human Hypoxia-inducible factor 1 alpha. To enhance the immunological response, this peptide was coupled to carrier protein BSA.
<b>Other Names:</b>	HIF-1 alpha; HIF1 alpha; ARNT-interacting protein; Member of PAS protein 1; Basic-helix-loop-helix-PAS protein MOP1; HIF1A; MOP1;
<b>Accession:</b>	Q16665 HIF1A_HUMAN;
<b>Produced in:</b>	Rabbit
<b>Purity:</b>	Affinity purified on antigen column
<b>Applications:</b>	Western Blotting (WB). A concentration of 2.0 µg/ml is recommended for WB. Human HIF-1 alpha (isoform 1) has a predicted length of 826 residues and a MW of 93 kDa. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	The specificity of this antibody has been confirmed by WB against the antigen.
<b>Cross-reactivity:</b>	Human; mouse; rat;
<b>Form:</b>	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg NaN <sub>3</sub>
<b>Reconstitution:</b>	Reconstitute in 100 µl of sterile distilled water to achieve an antibody concentration of 1 mg/ml. Centrifuge to remove any insoluble material.
<b>Storage:</b>	At least 12 months after purchase at 2 - 4°C (lyophilized formulations). After reconstitution, aliquot and store at -20°C for a higher stability. Avoid freeze-thaw cycles
<b>Expiry Date:</b>	12 months after purchase

---

FOR RESEARCH USE ONLY

## Rabbit polyclonal antibody to Hypoxia-inducible factor-1 alpha (775-790): Affinity purified



Western Blot using Rabbit polyclonal antibody to Hypoxia-inducible factor-1 alpha (R-1061-100). Lane 1: MCF-7 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: smmc whole cell lysate Lane 5: HT1080 whole cell lysate

biosensis

---

FOR RESEARCH USE ONLY