



Rabbit polyclonal antibody to Protein kinase C gamma (665-685): Affinity purified

Catalogue No.:	R-1142-100
Description:	Protein kinase C gamma (PKC-gamma) belongs to the Protein kinase C family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. These kinases phosphorylate a wide range of cellular proteins and are known to be involved in diverse cellular signaling pathways. PKC-gamma is one of at least 12 different types of Protein kinase C identified in humans. PKC-gamma is expressed in neurons of brain and spinal cord.
Batch No.:	See product label
Unit size:	100 ug
Antigen:	A synthetic peptide (IDQADFQGFTYVNPDFVHPDA) corresponding to a region (665-685) from rat Protein kinase C gamma. To enhance the immunological response, this peptide was coupled to carrier protein BSA.
Other Names:	PKC-gamma; EC 2.7.11.13; Prkcg; Pkcc; Pkcg; Prkcc;
Accession:	P63319 KPCG_RAT;
Produced in:	Rabbit
Purity:	Affinity purified on antigen column
Applications:	Western Blotting (WB). A concentration of 1.0 ug/mL is recommended for WB. Rat PKC-gamma has a predicted length of 697 residues and MW of 78 kDa. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The specificity of this antibody has been confirmed by WB against the antigen.
Cross-reactivity:	Rat; predicted to react with mouse due to sequence homology;
Form:	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃
Reconstitution:	Reconstitute in 100 uL of sterile distilled water to achieve an antibody concentration of 1 mg/mL. Centrifuge to remove any insoluble material.
Storage:	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
Expiry Date:	12 months after purchase

FOR RESEARCH USE ONLY