

## Mouse monoclonal antibody to Vascular endothelial growth factor receptor 2 [FLK-6]: IgG

<b>Catalogue No.:</b>	M-1183-100
<b>Description:</b>	Vascular endothelial growth factor (VEGF) receptor 2 is a type III receptor tyrosine kinase. As a receptor for VEGF, it mediates VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting.
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 ug
<b>Antigen:</b>	Recombinant human extracellular domain of Vascular endothelial growth factor receptor 2
<b>Other Names:</b>	VEGFR-2; Kinase insert domain receptor; Protein-tyrosine kinase receptor Flk-1; CD309; KDR; FLK1;
<b>Accession:</b>	P35968 VGFR2_HUMAN;
<b>Produced in:</b>	Mouse
<b>Purity:</b>	IgG
<b>Applications:</b>	Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 0.5-1.0 ug/mL is recommended for WB. Human VEGF receptor 2 has a predicted length of 1,356 residues and MW of 152 kDa. A concentration of 1.0-2.0 ug/mL is recommended to detect the protein in formalin/acetone fixed frozen tissues.
<b>Specificity:</b>	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
<b>Cross-reactivity:</b>	Human;
<b>Form:</b>	Lyophilized from 1.2% sodium acetate, 2mg BSA, 0.01mg NaN3
<b>Reconstitution:</b>	Reconstitute in 1 mL of PBS (pH 7.4) to achieve an antibody concentration of 100 ug/mL. Centrifuge to remove any insoluble material.
<b>Storage:</b>	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles
<b>Expiry Date:</b>	12 months after purchase

---

FOR RESEARCH USE ONLY

---