



## Mouse monoclonal antibody to alpha-Tubulin [Tub-1]: IgG

<b>Catalogue No.:</b>	M-1192-100
<b>Description:</b>	The tubulin family includes five distinct tubulins; alpha, beta, gamma, delta and epsilon. Microtubules consist principally of alpha- and beta-tubulin.
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 ug
<b>Antigen:</b>	Microtubules from chicken embryo brain
<b>Clone:</b>	Tub-1
<b>Other Names:</b>	Alpha-tubulin; Tubulin K-alpha-1; TUBA1B; TUBA1A; TUBA1C; TUBA3C; TUBA3D; TUBA3E; TUBA4A; TUBULIN; ALPHA-1A; Tubulin alpha-1B chain; Tubulin alpha-ubiquitous chain;
<b>Accession:</b>	P68363 TBA1B_HUMAN; P02552 TBA1_CHICK;
<b>Produced in:</b>	Mouse
<b>Purity:</b>	IgG
<b>Applications:</b>	Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 2.0 ug/mL is recommended for WB. Human Tubulin alpha-1B chain has a predicted length of 451 residues and MW of 50 kDa. A concentration of 1.0-2.0 ug/mL is recommended for IHC to detect the protein in formalin fixed and paraffin embedded tissues. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
<b>Cross-reactivity:</b>	Human; rat;
<b>Form:</b>	Lyophilized from 1.2% sodium acetate, 2mg BSA, 0.01mg NaN <sub>3</sub>
<b>Reconstitution:</b>	Reconstitute in 1 mL of PBS (pH 7.4) to achieve an antibody concentration of 100 ug/mL. Centrifuge to remove any insoluble material.
<b>Storage:</b>	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
<b>Expiry Date:</b>	12 months after purchase
<b>Specific References:</b>	Oh H. et al (2013) Uncovering a role for endocannabinoid signaling in autophagy in preimplantation mouse embryos. Mol Hum Reprod. 2013 Feb;19(2):93-101.

---

FOR RESEARCH USE ONLY