


## Rabbit polyclonal antibody to Tumor necrosis factor, alpha-induced protein 1, endothelial (27-52): Affinity purified

<b>Catalogue No.:</b>	R-1347-100
<b>Description:</b>	TNFAIP1 is a component of the BCR(BACURD2) E3 ubiquitin ligase complex (Ref: SWISSPROT).
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 ug
<b>Antigen:</b>	A synthetic peptide (NKYVQLNVGGSLYYTTVRALTRHDTM) corresponding to a region (27-52 aa) from the N-terminus of human tumor necrosis factor, alpha-induced protein 1, endothelial (TNFAIP1).
<b>Sequence:</b>	NKYVQLNVGGSLYYTTVRALTRHDTM
<b>Antibody Type:</b>	Antiserum
<b>Other Names:</b>	BTB/POZ domain-containing adapter for CUL3-mediated RhoA degradation protein 2; hBACURD2; BTB/POZ domain-containing protein TNFAIP1; Protein B12; TNFAIP1; BACURD2; EDP1;
<b>Accession:</b>	Q13829 BACD2_HUMAN;
<b>Produced in:</b>	Rabbit
<b>Applications:</b>	Immunohistochemistry (IHC), Immunocytochemistry (ICC) and Western Blotting (WB). A concentration of 0.5-1.0 ug/mL is recommended for WB. Human TNFAIP1 has a predicted length of 316 residues and MW of 36 kDa. A concentration of 0.25-0.5 ug/mL is recommended for IHC to detect the protein in formalin fixed and paraffin embedded tissues. Heat mediated antigen retrieval is required. A concentration of 0.25-0.5 ug/mL is recommended for IHC to detect the protein in frozen tissues. A concentration of 0.25-0.5 ug/mL is recommended for ICC. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
<b>Antibody Against:</b>	Tumor necrosis factor, alpha-induced protein 1, endothelial
<b>Cross-reactivity:</b>	Human; rat; predicted to react with mouse due to sequence homology;
<b>Form:</b>	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg Na <sub>3</sub>
<b>Appearance:</b>	Lyophilized white powder
<b>Reconstitution:</b>	Reconstitute in 100 uL of sterile distilled water to achieve an antibody concentration of 1 mg/mL. Centrifuge to remove any insoluble material.
<b>Storage:</b>	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles
<b>Expiry Date:</b>	12 months after purchase

---

FOR RESEARCH USE ONLY

## Rabbit polyclonal antibody to Tumor necrosis factor, alpha-induced protein 1, endothelial (27-52): Affinity purified



Western Blot using Rabbit polyclonal antibody to Tumor necrosis factor, alpha-induced protein 1, endothelial: Lane 1: Rat brain tissue lysate Lane 2: Rat Thymus tissue lysate Lane 3: Rat heart tissue lysate Lane 4: Rat lung tissue lysate

biosensis

---

FOR RESEARCH USE ONLY

---