



Chicken polyclonal antibody to Myelin Basic Protein (MBP): Affinity purified

Catalogue No.:	C-1383-50
Description:	Myelin is a membrane characteristic of the nervous tissue and functions as an insulator to increase the velocity of the stimuli being transmitted between a nerve cell body and its target. Myelin isolated from human and bovine nervous tissue is composed of approximately 80% lipid and 20% protein, and 30% of the protein fraction constitutes myelin basic protein (MBP). MBP is an intrinsically unstructured protein with a high proportion (approximately 75%) of random coil, but postulated to have core elements of beta-sheet and alpha-helix. MBP is a major protein in CNS myelin and is expressed specifically in the nervous system. A detailed immunochemical examination of monoclonal and polyclonal antibody responses to MBP and its peptides has revealed the existence of as many as 27 antigenic determinants, many of them conformational. Topological mapping of the potential antigenic determinants onto a model of MBP secondary structure places these determinants within 11 separate regions of the molecule, including those portions that have been found to be encephalitogenic. The message for myelin basic protein is selectively translocated to the ends of the cell processes. Immunization with myelin-associated antigens including MBP significantly promotes recovery after spinal cord contusion injury in the rat model. FUNCTION: Is, with PLP, the most abundant protein component of the myelin membrane in the CNS. Has a role in both the formation and stabilization of this compact multilayer arrangement of bilayers. Each splice variant and charge isomer may have a specialized function in the assembly of an optimized, biochemically functional myelin membrane (By similarity). SUBUNIT: Homodimer (By similarity). SUBCELLULAR LOCATION: Myelin membrane; peripheral membrane protein; cytoplasmic side. Cytoplasmic side of myelin. TISSUE SPECIFICITY: Found in both the central and the peripheral nervous system.
Batch No.:	See product label
Unit size:	50 uL
Antigen:	Three peptide sequences conserved in higher vertebrate MBP protein.
Isotype:	IgY
Other Names:	Myelin Basic Protein; Myelin A1 protein; Myelin membrane encephalitogenic protein;
Accession:	P02686 MBP_HUMAN;
Produced in:	Chicken
Applications:	Western Blotting (WB), Immunocytochemistry (IC) and Immunohistochemistry (IH). The recommended dilution for WB is 1:5,000-10,000 and 1:500-1,000 for IC and IH. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The specificity of this antibody has been confirmed by WB. This antibody stains a prominent band at approx. 20 kDa. A suitable control tissue is rat spinal cord or peripheral nerve homogenate. The major isoforms of MBP run as a closely spaced double of 22 kDa and 18 kDa.
Antibody Against:	Myelin Basic Protein

FOR RESEARCH USE ONLY



Chicken polyclonal antibody to Myelin Basic Protein (MBP): Affinity purified

- Cross-reactivity:** Hu, Rat, Ms, Bov. Predicted to react with other mammalian tissues due to sequence homology.
- Form:** Lyophilised with 5% trehalose
- Appearance:** White powder
- Reconstitution:** Reconstitute in sterile distilled water. Centrifuge to remove any insoluble material.
- Storage:** After reconstitution of lyophilised antibody, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
- Expiry Date:** 12 months after purchase
- Specific References:** 1. Rangaraju S. et al (2009) Molecular architecture of myelinated peripheral nerves is supported by calorie restriction with aging. *Aging Cell*. 2009 Apr;8(2):178-91.




Image shows rat mixed neuron/glia cultures stained with Chicken polyclonal antibody to Myelin Basic Protein antibody C-1383-50 (green). Blue is a DNA stain. Note that the antibody stains an oligodendrocyte and some membrane shed from this cell. Other cells in the field include neurons, astrocytes, microglia and fibroblasts, all of which are completely negative.

FOR RESEARCH USE ONLY