

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Catalogue No.: TR-100-FJT

Description: The causes and effects of neuronal degeneration are of major interest to a wide variety of neuroscientists. Paralleling this growing interest is an increasing number of methods applicable to the detection of neuronal degeneration. Fluoro-Jade C stains all degenerating neurons regardless of specific insult or mechanism of cell death. Fluoro-Jade C exhibits the greatest signal to background ratio, as well as the highest resolution. This translates to a stain of maximal contrast and affinity for degenerating neurons. This makes it ideal for localising not only degenerating nerve cell bodies but also distal dendrites, axons and terminals. The dye is highly resistant to fading and is compatible with virtually all histological processing and staining protocols.

Applications: The Fluoro-Jade C 'Ready to Dilute' (RTD) Staining Kit provides an easy to use assortment of Fluoro-Jade C, DAPI, sodium hydroxide and potassium permanganate in liquid form. Following our detailed protocol, Fluoro-Jade C labelled degenerating neurons are visualised with blue light excitation while DAPI counter stained cell nuclei are visualised with ultra-violet illumination. The Fluoro-Jade C Staining Kit can be used on all kinds of preserved tissues, including fresh-frozen, paraformaldehyde or formalin fixed, and formalin fixed, paraffin-embedded tissues.

Comments: MATERIALS PROVIDED

Sodium Hydroxide, Solution A (Dilute 1:10 prior to use) - 20 mL
Potassium Permanganate, Solution B (Dilute 1:10 prior to use) - 20 mL
Fluoro-Jade C, Solution C (Dilute 1:10 prior to use) - 20 mL
DAPI, Solution D (Add to diluted Fluoro-Jade C) - 20 mL

EQUIPMENT AND REAGENTS NEEDED

Gelatin coated microscope slides
Staining dishes/Coplin jars
Cover slips
DPX mounting media
Slide warmer
Convection oven
Distilled water
Ethanol
Xylene

Final working concentrations of FJC:0.0001%
Final working concentration of KMnO₄: 0.06%
NUMBER OF SLIDES PROCESSED:

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

The actual number of slides processed by this kit will depend largely upon the vessel that is used to incubate the slides. If using a standard Coplin Jar, its capacity is 50 mL and typically holds 5 slides per jar. If using such a device, then 80-100 slides stained per 50 ml of working solution (or, 5 ml of stock solution) could be processed in one day. Note the diluted dye is NOT stable and will not store overnight. It is best to use freshly diluted dye each time an experimental batch is started.

Specificity: Degenerating neurons, and neuronal degeneration. There is no specific staining in normal healthy brain.

Cross-reactivity: Note: Some researchers under some conditions report blood vessel staining with Fluoro Jade. This may be because Fluoro Jade is an analogue of eosin (which stains blood cells). In general, good perfusion and preparation of the tissue should help prevent blood vessel staining but it may not be possible to eliminate it entirely. In our experience it is generally possible to distinguish neuronal from blood vessels staining by eye.

Form: The reagents in the Fluoro Jade kit are all supplied in a liquid format and are ready-to-dilute.

Appearance: FJC visualization is accomplished using blue light or a 488nm Laser.

Excitation Peak: 495 nm

Emission Peak: 521 nm

Filter system for visualizing: Fluorescein/FITC

Reconstitution: Dilute solutions as directed in the protocol instructions. Sometimes small precipitates may be present in the stock or diluted solutions. Complete mixing of the diluted solutions usually dissolves the precipitates. The precipitates, if not removed, do not usually cause any difficulties if the washing steps are followed as instructed. Optional: For entirely clean solutions Biosensis recommends filtering the diluted solution through ethanol and NaOH compatible syringe or vacuum filter devices prior to contact with tissue slides.

Storage: The unopened kit can be stored for up to 6 months at 2-8C after the date of receipt. The kit and components should be stored protected from light. Diluted FJC dye solutions are not stable and should be used within 4 hours of making. The other diluted solutions can be reused and stored for up to 48 hours if refrigerated and protected from light. Best results require freshly diluted solutions. We recommend using aseptic techniques when handling the reagents to avoid bacterial growth and contamination. The FJC Ready to dilute kit is shipped ambient and stable at room temperature during transport. Refrigerate upon arrival, do not freeze.

Expiry Date: Unopened kit 6 months at 2-8C protected from light. See Storage instructions for working solutions recommendations.

Specific References: Hu H et al. (2020) "Transient receptor potential melastatin 2 contributes to neuroinflammation and negatively regulates cognitive outcomes in a pilocarpine-induced mouse model of epilepsy." *Int Immunopharmacol.* 87:106824. Application: IF.

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Wang S et al. (2020) "Aging exacerbates impairments of cerebral blood flow autoregulation and cognition in diabetic rats." *Geroscience*. [Epub ahead of print]. Application: IF.

Chen PY et al. (2020) "Stearic Acid Methyl Ester Affords Neuroprotection and Improves Functional Outcomes after Cardiac Arrest." *Prostag Leukotr Ess*. [In Press]. Application: IF.

Gamdzyk M et al. (2020) "cGAS/STING Pathway Activation Contributes to Delayed Neurodegeneration in Neonatal Hypoxia-Ischemia Rat Model: Possible Involvement of LINE-1." *Mol Neurobiol*. 57(6): 2600-19. Application: IF. Species: Rat

Tang H et al. (2020) "Delayed Recanalization After MCAO Ameliorates Ischemic Stroke by Inhibiting Apoptosis via HGF/c-met/STAT3/Bcl-2 Pathway in Rats." *Exp Neurol*. [Epub ahead of print]. Application: IF. Species: Rat

Ocak U et al. (2020) "Inhibition of mast cell tryptase attenuates neuroinflammation via PAR-2/p38/NF- κ B pathway following asphyxial cardiac arrest in rats." *J Neuroinflammation*. 17(1):144. Application: IF. Species: Rat

Huang J et al. (2020) "IRE1 α ; inhibition attenuates neuronal pyroptosis via miR-125/NLRP1 pathway in a neonatal hypoxic-ischemic encephalopathy rat model." *J Neuroinflammation*. 17(1):152. Application: IF. Species: Rat

Fouda MA et al. (2020) "Estrogen-dependent hypersensitivity to diabetes-evoked cardiac autonomic dysregulation: Role of hypothalamic neuroinflammation." *Life Sci*. 2020 Mar 31:117598. [Epub ahead of print]. Application: IF. Species: Rat

Zhou Z et al. (2020) "Sodium butyrate attenuated neuronal apoptosis via GPR41/G β 17;PI3K/Akt pathway after MCAO in rats." *J Cerebr Blood F Met*. [Epub ahead of print]. Application: IF. Species: Rat

Enam SF et al. (2020) "Engineering Cytokine and Macrophage Enrichment at Sites of Injury." PhD Thesis. Application: IF. Species: Mouse

Lynch-Godrei A et al. (2019) "Characterization of gastrointestinal pathologies in the dystonia musculorum mouse model for hereditary sensory and autonomic neuropathy type VI." *Neurogastroenterol Motil*. [Epub ahead of print]. Application: IF. Species: Mouse

Ito T et al. (2019) "Vestibular Compensation after Vestibular Dysfunction Induced by Arsanilic Acid in Mice." *Brain Sci*. 9(11):329. Application: IF. Species: Mouse

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Miao H et al. (2019) "Edaravone Administration Confers Neuroprotection after Experimental Intracerebral Hemorrhage in Rats via NLRP3 Suppression." *J Stroke Cerebrovasc Dis.* [Epub ahead of print]. Application: IF. Species: Rat

Ocak U J et al. (2019) "FSLRY-NH₂ Improves Neurological Outcome After Cardiac Arrest in Rats." *Turk Neurosurg.* Application: IF. Species: Rat

Huang J et al. (2019) "Ghrelin attenuates oxidative stress and neuronal apoptosis via GHSR-1α/AMPK/Sirt1/PGC-1α/UCP2 pathway in a rat model of neonatal HIE." *Free Radic Biol Med.* 141:322-37. Application: IF. Species: Rat

Sun C et al. (2019) "Osteopontin-Enhanced Autophagy Attenuates Early Brain Injury via FAK-ERK Pathway and Improves Long-Term Outcome after Subarachnoid Hemorrhage in Rats." *Cells.* [Epub ahead of print]. Application: IF. Species: Rat

Zuo G et al. (2019) "Activation of TGR5 with INT-777 attenuates oxidative stress and neuronal apoptosis via cAMP/PKCε/ALDH2 pathway after subarachnoid hemorrhage in rats." *Free Radic Biol Med.* [Epub ahead of print]. Application: IF. Species: Rat

Carron SF et al. (2019) "Inhibitory neuronal changes following a mixed diffuse-focal model of traumatic brain injury." *J Comp Neurol.* [Epub ahead of print]. Application: IF. Species: Rat

Sepulveda-Rodriguez A (2019) "Acute Microglial Responses to Single Non-Epileptogenic vs. Epileptogenic Seizures in Mouse Hippocampus." PhD Thesis. Application: IF. Species: Mouse

Nomura E et al. (2019) "Imaging Hypoxic Stress and the Treatment of Amyotrophic Lateral Sclerosis with Dimethylxalylglycine in a Mice Model." *Neuroscience.* [Epub ahead of print]. Application: IF. Species: Mouse

Okada T et al. (2019) "FGF-2 Attenuates Neuronal Apoptosis via FGFR3/PI3k/Akt Signaling Pathway After Subarachnoid Hemorrhage." *Mol Neurobiol.* [Epub ahead of print]. Application: IF. Species: Rat

Sepulveda-Rodriguez A et al. (2019) "Electroconvulsive shock enhances responsive motility and purinergic currents in microglia in the mouse hippocampus." *eNeuro.* Application: IF. Species: Mouse

Teng L et al. (2019) "Carnosic Acid Mitigates Early Brain Injury After Subarachnoid Hemorrhage: Possible Involvement of the SIRT1/p66shc Signaling Pathway." *Front Neurosci.* 13:26. Application: IF. Species: Rat

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Chen S et al. (2019) "RO27-3225 attenuates NLRP1-dependent neuronal pyroptosis via MC4R/ASK1/JNK/p38 MAPK pathway in a mouse model of intracerebral hemorrhage." *Br J Pharmacol.* [Epub ahead of print]. Application: IF. Species: Mouse

Batsaikhan B et al. (2019) "Post-Injury Neuroprotective Effects of the Thalidomide Analog 3,6-dithiothalidomide on Traumatic Brain Injury." *Int J Mol Sci.* 20(3). Application: IF. Species: Rat

Dominy SS et al. (2019) "Porphyromonas gingivalis in Alzheimer's disease brains: Evidence for disease causation and treatment with small-molecule inhibitors." *Sci Adv.* 5(1). Application: IF. Species: Mouse

Zhao L et al. (2019) "Administration of rCTRP9 Attenuates Neuronal Apoptosis Through AdipoR1/PI3K/Akt Signaling Pathway after ICH in Mice." *Cell Transplant.* [Epub ahead of print]. Application: IF. Species: Mouse

Mao Y et al. (2018) "The Effects of a Combination of Ion Channel Inhibitors in Female Rats Following Repeated Mild Traumatic Brain Injury." *Int J Mol Sci.* 19(11). Application: IF. Species: Rat

Yon JM et al. (2018) "The Ethanol Fraction of White Rose Petal Extract Abrogates Excitotoxicity-Induced Neuronal Damage In Vivo and In Vitro through Inhibition of Oxidative Stress and Proinflammation." *Nutrients.* 10(10). Application: IF. Species: Mouse

Yu J et al. (2018) "Ezetimibe, a NPC1L1 inhibitor, attenuates neuronal apoptosis through AMPK dependent autophagy activation after MCAO in rats." *Exp Neurol.* 307:12-23. Application: IF. Species: Rat

Nguyen TV et al. (2018) "Alzheimer's associated amyloid and tau deposition co-localizes with a homeostatic myelin repair pathway in two mouse models of post-stroke mixed dementia." *Acta Neuropathol Commun.* 6(1):100. Application: IF. Species: Mouse

Mo J et al. (2018) "AVE 0991 attenuates oxidative stress and neuronal apoptosis via Mas/PKA/CREB/UCP-2 pathway after subarachnoid hemorrhage in rats." *Redox Biol.* 20:75-86. Application: IF. Species: Rat

Nito C et al. (2018) "Transplantation of human dental pulp stem cells ameliorates brain damage following acute cerebral ischemia." *Biomed Pharmacother.* 108:1005-14. Application: IF. Species: Rat

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Mamo JC et al. (2018) "Sodium alginate capsulation increased brain delivery of probucol and suppressed neuroinflammation and neurodegeneration." *Ther Deliv.* 9(10):703-9. Application: IF. Species: Mouse

Ilievski V et al. (2018) "Chronic oral application of a periodontal pathogen results in brain inflammation, neurodegeneration and amyloid beta production in wild type mice." *PLoS One.* 13(10):e0204941. Application: IF. Species: Mouse

Mamo JC et al. (2018) "Probucol prevents blood-brain barrier dysfunction and cognitive decline in mice maintained on pro-diabetic diet." *Diab Vasc Dis Res.* [Epub ahead of print]. Application: IF. Species: Mouse

Fang C et al. (2018) "Tanshinone IIA improves hypoxic ischemic encephalopathy through TLR4-mediated NF- κ B signal pathway." *Mol Med Rep.* 18(2):1899-1908. Application: IF. Species: Mouse

Hubbard WB et al. (2018) "Hemostatic nanoparticles increase survival, mitigate neuropathology and alleviate anxiety in a rodent blast trauma model." *Sci Rep.* 8(1):10622. Application: IF. Species: Rat

Sowa K et al. (2018) "Impact of Dental Pulp Stem Cells Overexpressing Hepatocyte Growth Factor after Cerebral Ischemia/Reperfusion in Rats." *Mol Ther Methods Clin Dev.* [Epub ahead of print]. Application: IF. Species: Rat

Nakajima M, Suda S, Sowa K, Sakamoto Y, Nito C, Nishiyama Y, Aoki J, Ueda M, Yokobori S, Yamada M, Yokota H, Okada T, Kimura K (2018) "AMPA receptor antagonist perampanel ameliorates post-stroke functional and cognitive impairments." *Neuroscience.* [Epub ahead of print]. Application: IF. Species: Rat

Matei N, Camara J, McBride D, Camara R, Xu N, Tang J, Zhang JH (2018) "Intranasal wnt3a attenuates neuronal apoptosis through Frz1/PIWIL1a/FOXO1 pathway in MCAO rats." *J Neurosci.* [Epub ahead of print]. Application: IF. Species: Rat

Ignowski E, Winter AN, Duval N, Fleming H, Wallace T, Manning E, Koza L, Huber K, Serkova NJ, Linseman DA (2018) "The cysteine-rich whey protein supplement, Immunocal®, preserves brain glutathione and improves cognitive, motor, and histopathological indices of traumatic brain injury in a mouse model of controlled cortical impact." *Free Radic Biol Med.* [Epub ahead of print]. Application: IF. Species: Mouse

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Shao Z, Wu P, Wang X, Jin M, Liu S, Ma X, Shi H (2018) "Tetramethylpyrazine Protects Against Early Brain Injury and Inhibits the PERK/Akt Pathway in a Rat Model of Subarachnoid Hemorrhage." *Neurochem Res.* [Epub ahead of print]. Application: IF. Species: Rat

Koyama K, Kangawa A, Fukumoto N, Watanabe KI, Horiuchi N, Ozawa T, Inokuma H, Kobayashi Y (2018) "Histopathological study of encephalomalacia in neonatal calves and application of neuronal and axonal degeneration marker." *J Vet Med Sci.* [Epub ahead of print]. Application: IF. Species: Cow

Fouda MA, El-Sayed SS, Abdel-Rahman AA (2018) "Restoration of Rostral Ventrolateral Medulla Cystathionine- γ -Lyase Activity Underlies Moxonidine-Evoked Neuroprotection and Sympathoinhibition in Diabetic Rats." *J Pharmacol Exp Ther.* 364(2):170-8 Application: IF. Species: Rat

Huang YN, Yang LY, Greig NH, Wang YC, Lai CC, Wang JY (2018) "Neuroprotective effects of pifithrin- α ; against traumatic brain injury in the striatum through suppression of neuroinflammation, oxidative stress, autophagy, and apoptosis." *Sci Rep.* 8(1): 2368 Application: IF. Species: Rat

Takechi R, Lam V, Brook E, Giles C, Fimognari N, Mooranian A, Al-Salami H, Coulson SH, Nesbit M, Mamo JCL (2017) "Blood-Brain Barrier Dysfunction Precedes Cognitive Decline and Neurodegeneration in Diabetic Insulin Resistant Mouse Model: An Implication for Causal Link." *Front Aging Neurosci.* 9:399 Application: IF. Species: Mouse

Freitas-Andrade M, She J, Bechberger J, Naus CC, Sin WC (2017) "Acute connexin43 temporal and spatial expression in response to ischemic stroke." *J Cell Commun Signal.* [Epub ahead of print] Application: IF. Species: Mouse

Xie Z, Enkhjargal B, Wu L, Zhou K, Sun C, Hu X, Gospodarev V, Tang J, You C, Zhang JH (2017) "Exendin-4 attenuates neuronal death via GLP-1R/PI3K/Akt pathway in early brain injury after subarachnoid hemorrhage in rats." *Neuropharmacology.* 128:142-51 Application: IF. Species: Rat

Khan D (2017) "The Relationship between Astrocytes, Inflammation and Epileptogenesis." PhD Thesis. Application: IF. Species: Mouse

Fan LF, He PY, Peng YC, Du QH, Ma YJ, Jin JX, Xu HZ, Li JR, Wang ZJ, Cao SL, Li T, Yan F, Gu C, Wang L, Chen G (2017) "Mdivi-1 ameliorates early brain injury after subarachnoid hemorrhage via the suppression of inflammation-related blood-brain barrier disruption and endoplasmic reticulum stress-based apoptosis." *Free Radic Biol Med.* [Epub ahead of print]

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Application: IF. Species: Rat

Moro C, Torres N, Arvanitakis K, Cullen K, Chabrol C, Agay D, Darlot F, Benabid AL, Mitrofanis J (2017) "No evidence for toxicity after long-term photobiomodulation in normal non-human primates." *Exp Brain Res*. [Epub ahead of print] Application: IF. Species: Primate

Fan R, Enkhjargal B, Camara R, Yan F, Gong L, ShengtaoYao, Tang J, Chen Y, Zhang JH (2017) "Critical role of EphA4 in early brain injury after subarachnoid hemorrhage in rat." *Exp Neurol*. 296:41-8 Application: IF. Species: Rat

Nakajima M, Nito C, Sowa K, Suda S, Nishiyama Y, Nakamura-Takahashi A, Nitahara-Kasahara Y, Imagawa K, Hirato T, Ueda M, Kimura K, Okada T (2017) "Mesenchymal Stem Cells Overexpressing Interleukin-10 Promote Neuroprotection in Experimental Acute Ischemic Stroke." *Mol Ther Methods Clin Dev*. 6:102-11 Application: IF. Species: Human

Suzuki K, Yamada K, Fukuhara Y, Tsuji A, Shibata K, Wakamatsu N (2017) "High-dose thiamine prevents brain lesions and prolongs survival of Slc19a3-deficient mice." *PLoS One*. 12(6):e0180279 Application: IF. Species: Mouse

Kononenko NL, Claßen GA, Kuijpers M, Puchkov D, Maritzen T, Tempes A, Malik AR, Skalecka A, Bera S, Jaworski J, Haucke V (2017) "Retrograde transport of TrkB-containing autophagosomes via the adaptor AP-2 mediates neuronal complexity and prevents neurodegeneration." *Nat Commun*. 8:14819 Application: IF. Species: Mouse

Lian YD, Chen ZX, Zhu KR, Sun SY, Zhu LP (2017) "Effect of equipotent doses of bupivacaine and ropivacaine in high-fat diet fed neonatal rodent model." *Braz J Anesthesiol*. 67(2):131-8 Application: IF. Species: Rat

Mamo JCL, Lam V, Giles C, Coulson SH, Fimognari N, Mooranian A, Al-Salami H, Takechi R (2017) "Antihypertensive agents do not prevent blood-brain barrier dysfunction and cognitive deficits in dietary-induced obese mice." *Int J Obes (Lond)*. [Epub ahead of print] Application: IF. Species: Mouse

Iwasa K, Yamamoto S, Yagishita S, Maruyama K, Yoshikawa K (2017) "Excitotoxicity-induced prostaglandin D2 production induces sustained microglial activation and delayed neuronal death." *J Lipid Res*. 58(4):649-55 Application: IF. Species: Rat

Xie Z, Huang L, Enkhjargal B, Reis C, Wan W, Tang J, Cheng Y, Zhang JH (2017) "Intranasal administration of recombinant Netrin-1 attenuates neuronal apoptosis by activating DCC/APPL-1/AKT signaling pathway after subarachnoid hemorrhage in rats."

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Neuropharmacology. <http://dx.doi.org/10.1016/j.neuropharm.2017.03.025> [In press] Application: IF. Species: Rat

Katayama Y, Inaba T, Nito C, Suda S, Ueda M (2016) "Neuroprotective effects of clarithromycin against neuronal damage in cerebral ischemia and in cultured neuronal cells after oxygen-glucose deprivation." *Life Sci.* PMID:27825902 [Epub ahead of print] Application: IF. Species: Rat

Yang LY, Greig NH, Huang YN, Hsieh TH, Tweedie D, Yu QS, Hoffer BJ, Luo Y, Kao YC, Wang JY (2016) "Post-traumatic administration of the p53 inactivator pifithrin- α ; oxygen analogue reduces hippocampal neuronal loss and improves cognitive deficits after experimental traumatic brain injury." *Neurobiol Dis.* PMID:27553877 [Epub ahead of print] Application: IF. Species: Rat

Izumida H, Takagi H, Fujisawa H, Iwata N, Nakashima K, Takeuchi S, Iwama S, Namba T, Komatu Y, Kaibuchi K, Oiso Y, Arima H, Sugimur Y (2016) "NMDA receptor antagonist prevents cell death in the hippocampal dentate gyrus induced by hyponatremia accompanying adrenal insufficiency in rats." *Exp Neurol.* PMID:27527984 [Epub ahead of print] Application: IF. Species: Rat

Dixon BJ, Chen D, Zhang Y, Flores J, Malaguit J, Nowrangi D, Zhang JH, Tang J. (2016) "Intranasal Administration of Interferon Beta Attenuates Neuronal Apoptosis via the JAK1/STAT3/BCL-2 Pathway in a Rat Model of Neonatal Hypoxic-Ischemic Encephalopathy." *ASN Neuro.* PMID:27683877 8(5) Application: IF. Species: Rat

Wang JY, Huang YN, Chiu CC, Tweedie D, Luo W, Pick CG, Chou SY, Luo Y, Hoffer BJ, Greig NH, Wang JY (2016) "Pomalidomide mitigates neuronal loss, neuroinflammation, and behavioral impairments induced by traumatic brain injury in rat." *J Neuroinflammation.* PMID:27353053 13(1):168 Application: IF. Species: Rat

Wang G, Manaenko A, Shao A, Ou Y, Yang P, Budbazar E, Nowrangi D, Zhang JH, Tang J (2016) "Low-density lipoprotein receptor-related protein-1 facilitates heme scavenging after intracerebral hemorrhage in mice." *J Cereb Blood Flow Metab.* PMID:27317656 [Epub ahead of print] Application: IF. Species: Mouse

Safdaria BK, Siab TC, Wattchow DA, Smid SD (2016) "Effects of pro-inflammatory cytokines, lipopolysaccharide and COX-2 mediators on human colonic neuromuscular function and epithelial permeability." *Cytokine.* PMID:27177092 83:231-8 Application: IF. Species: Human

Lian YD, Chen ZX, Zhu KR, Sun SY, Zhu LP (2016) "Effect of equipotent doses of bupivacaine

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

and ropivacaine in high-fat diet fed neonatal rodent model." Brazilian Journal of Anesthesiology [In Press] Application: IF. Species: Rat

Launay PS, Reboussin E, Liang H, Kessal K, Godefroy D, Rostene W, Sahel JA, Baudouin C, Parsadaniantz SM, Le Goazigo AR (2016) "Ocular inflammation induces trigeminal pain, peripheral and central neuroinflammatory mechanisms." Neurobiology of Disease. PMID:26747211 88:16-28 Application: IH. Species: Mouse

Khan D, Dupper A, Deshpande T, Graan PN, Steinhäuser C, Bedner P. (2016) "Experimental febrile seizures impair interastrocytic gap junction coupling in juvenile mice." J Neurosci Res. PMID:26931373 [Epub ahead of print] Application: IF. Species: Mouse

Bye N, Christie KJ, Turbic A, Basrai HS, Turnley AM (2016) "Rho kinase inhibition following traumatic brain injury in mice promotes functional improvement and acute neuron survival but has little effect on neurogenesis, glial responses or neuroinflammation." Exp Neurol. PMID:26896832 [Epub ahead of print] Application: IH. Species: Mouse

Bedner P, Dupper A, Hüttmann K, Müller J, Herde MK, Dublin P, Deshpande T, Schramm J, Häussler U, Haas CA, Henneberger C, Theis M, Steinhäuser C (2015) "Astrocyte uncoupling as a cause of human temporal lobe epilepsy." Brain. PMID:25765328 [Epub ahead of print] Application: IH 4%PFA, vibratome 40um sections. Species: Mouse

Ramesh G, Didier PJ, England JD, Santana-Gould L, Doyle-Meyers LA, Martin DS, Jacobs MB, Philipp MT (2015) "Inflammation in the Pathogenesis of Lyme Neuroborreliosis." Am J Pathol. PMID:25892509 [Epub ahead of print] Application: IH with Z-fixative, 5 micron sections. Species: Rhesus Monkey

Suda S, Ueda M, Nito C, Nishiyama Y, Okubo S, Abe A, Aoki J, Suzuki K, Sakamoto Y, Kimura K (2015) "Valproic acid ameliorates ischemic brain injury in hyperglycemic rats with permanent middle cerebral occlusion." Brain Res. PMID:25721785 1606:1-8 Application: IH. Species: Rat

Mayeux JP, Teng SX, Katz PS, Gilpin NW, Molina PE (2015) "Traumatic brain injury induces neuroinflammation and neuronal degeneration that is associated with escalated alcohol self-administration in rats." Behav Brain Res. PMID:25446758 279:22-30 Application: IH. Species: Rat

Yang LY, Chu YH, Tweedie D, Yu QS, Pick CG, Hoffer BJ, Greig NH, Wang JY (2015) "Post-trauma administration of the pifithrin- α ; oxygen analog improves histological and functional outcomes after experimental traumatic brain injury." Exp Neurol. PMID:25819102 269:56-66 Application: IH. Species: Rat

FOR RESEARCH USE ONLY

Fluoro-Jade C (FJC) Ready-to-Dilute Staining Kit for identifying Degenerating Neurons (Trial size)

Ren YY, Zhang HQ, Duan SH, Wang SD (2015) "Influence of Dexmedetomidine on Toxicity of Intrathecal

Ketamine on Neonatal Rat Spinal Function." *Tropical Journal of Pharmaceutical Research*. 14(3):469-77 Application: IH. Species: Rat


Iseli CE, Merwin WH 3rd, Klatt-Cromwell C, Hutson KA, Ewend MG, Adunka OF, Fitzpatrick DC, Buchman CA (2014) "Effect of Cochlear Nerve Electrocautery on the Adult Cochlear Nucleus." *Otol Neurotol*. PMID: 25280052 [Epub ahead of print] Application: IH(P)

Saito T, Nito C, Ueda M, Inaba T, Kamiya F, Muraga K, Katsura K, Katayama Y (2014) "Continuous oral administration of atorvastatin ameliorates brain damage after transient focal ischemia in rats." *Life Sci*. PMID: 24333133 94(2):106-14 Application: IH. Species: Rat

Hayashida K, Sano M, Kamimura N, Yokota T, Suzuki M, Ohta S, Fukuda K, Hori S (2014) "Hydrogen inhalation during normoxic resuscitation improves neurological outcome in a rat model of cardiac arrest independently of targeted temperature management." *Circulation*. PMID: 25366995 130(24):2173-80 Application: IH. Species: Rat

Reagent Kit protocol: Please refer to our online product listing for current protocol/MSDS versions.

MSDS: Please refer to our online product listing for current protocol/MSDS versions.



Fluoro-Jade C (FJC) staining (green) of degenerating neurons in the CA3 region of the hippocampus in old diabetic rats. Blue: DAPI nuclear stain. Picture courtesy of Dr. Wang and colleagues. Results of this study were published in Wang S. et al. (2020), *Geroscience*, PMID: 32696219.

FOR RESEARCH USE ONLY