

## Rabbit polyclonal antibody to human Glial Fibrillary Acidic Protein (341-356): Affinity purified

Catalogue No.: R-1145-100

Description: Glial Fibrillary Acidic Protein (GFAP) is a 50 kDa intra-cytoplasmic filamentous protein of the

cytoskeleton in astrocytes. During the development of the central nervous system, it is a cell-specific marker that distinguishes astrocytes from other glial cells. GFAP immunoreactivity has been shown in immature oligodendrocytes, epiglottic cartilage, pituicytes, papillary meningiomas, myoepithelial cells of the breast and in non-CNS: Schwann cells, salivary gland neoplasms, enteric glia cells, and metastasizing renal carcinomas. At least 3 isoforms are produced from alternate splicing. These isoforms differ in the C-terminal region which is

encoded by alternative exons.

Batch No.: See product label

Unit size: 100 ug

Antigen: A synthetic peptide (EMARHLQEYQDLLNVK) corresponding to a region (341-356) from human

Glial Fibrillary Acidic Protein. To enhance the immunological response, this peptide was

coupled to carrier protein BSA.

Other Names: Astrocyte; Glial fibrillary acidic protein; GFAP;

Accession: P14136 GFAP\_HUMAN;

Produced in: Rabbit

Purity: Affinity purified on antigen column

Applications: Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 1.0 ug/mL is

recommended for WB. Human GFAP has a predicted length of 432 residues and MW of 50 kDa. A concentration of 2.0 ug/mL is recommended to detect GFAP in formalin fixed and paraffin embedded tissues. Heat mediated antigen retrieval is required. Biosensis recommends

optimal dilutions/concentrations should be determined by the end user.

**Specificity:** The specificity of this antibody has been confirmed by WB and IHC against the antigen.

Cross-reactivity: Human; rat; predicted to react with mouse due to sequence homology;

Form: Lyophilized from 1.2% sodium acetate, 2mg BSA, 0.2mg NaN3

Reconstitution: Reconstitute in 1 mL of PBS (pH 7.4) to achieve an antibody concentration of 100 ug/mL.

Centrifuge to remove any insoluble material.

Storage: At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution,

aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.

**Expiry Date:** 12 months after purchase.

**References:** 1. Reeves S.A, et al. Proc. Natl. Acad. Sci. U.S.A. 86:5178-5182(1989).

2. Brenner M, et al. Brain Res. Mol. Brain Res. 7:277-286(1990).

Isaacs A, et al. Genomics 51:152-154(1998).
Ota T, et al. Nat. Genet. 36:40-45(2004).

4. Nielsen A.L, et al. J. Biol. Chem. 277:29983-29991(2002).

5. Singh R, et al. Genomics 82:185-193(2003).

## FOR RESEARCH USE ONLY



## Rabbit polyclonal antibody to human Glial Fibrillary Acidic Protein (341-356): Affinity purified

- 6. Brenner M, et al. Nat. Genet. 27:117-120(2001).
- 7. Brockmann K, et al. Eur. Neurol. 50:100-105(2003).
- 8. Stumpf E, et al. Arch. Neurol. 60:1307-1312(2003).
- 9. Sawaishi Y, et al. Neurology 58:1541-1543(2002).
- 10. Aoki Y, et al. Neurosci. Lett. 312:71-74(2001).

