


## Mouse monoclonal antibody to Neurofilament Light [DA2]

<b>Catalogue No.:</b>	M-1391-50
<b>Description:</b>	Neurofilaments are composed of three intermediate filament proteins: light (~68 kDa), medium (~160 kDa) and heavy (~200 kDa), which are involved in the maintenance of the neuronal caliber. Neurofilament light (NF68 or NF-L) is the most abundant of the three proteins.
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	50 uL
<b>Antigen:</b>	Enzymatically dephosphorylated full length pig NF-L protein. The antibody binding epitope has been mapped to a short peptide in the C-terminal "tail" region of the molecule within the sequence YYTSHVQEEQIEVEETIEA, amino acids 441-460 of the human sequence.
<b>Antibody Type:</b>	Monoclonal
<b>Isotype:</b>	IgG1
<b>Clone:</b>	DA2
<b>Other Names:</b>	NF-L; NF68; NEFL; Neurofilament light polypeptide; NFL;
<b>Accession:</b>	P02547 NFL_PIG; P07196 NFL_HUMAN;
<b>Produced in:</b>	Mouse
<b>Applications:</b>	Western Blotting (WB), Immunocytochemistry (IC), Immunohistochemistry (IH) and Flow Cytometry (2 ug per 10 <sup>6</sup> cells). A dilution of 1:5,000 - 1:10,000 is recommended for WB. A dilution of 1:100 - 1:500 is recommended for IC and IH. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	Specifically recognizes the light neurofilament subunit NF-L (~68 kDa) in WB.
<b>Antibody Against:</b>	Neurofilament Light
<b>Cross-reactivity:</b>	Hu, Rat, Ms, Fel, Bov, Por, Chk
<b>Form:</b>	Lyophilised with 5% trehalose
<b>Appearance:</b>	White powder
<b>Reconstitution:</b>	Reconstitute in sterile distilled water. Centrifuge to remove any insoluble material.
<b>Storage:</b>	After reconstitution of lyophilised antibody, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
<b>Expiry Date:</b>	12 months after purchase
<b>Specific References:</b>	1. Felitsyn N. et al (2008) The heme precursor delta-aminolevulinate blocks peripheral myelin formation. J Neurochem. 2008 Sep;106(5):2068-79.

---

FOR RESEARCH USE ONLY

## Mouse monoclonal antibody to Neurofilament Light [DA2]



Left: Cells grown from adult rat brain. The large cell in the middle is stained with Mouse monoclonal antibody to Neurofilament Light [DA2] M-1391-50 (green). Another type of neuronal lineage cell was stained with Rabbit polyclonal to alpha-internexin R-1379-50 (red). These cells were mitotic but had several characteristics of neurons. Right: Analysis of neurofilament, light, expression in human prostate cancer cell line DU145 by Flow Cytometry. Fixing and Permeabilization of cells: Absolute methanol (10 minutes in ice) and 0.1% Tween-20 in PBS, Blocking: 200 ug/mL Normal Sheep IgG (20 minutes), Primary antibody: Mouse Monoclonal antibody to Neurofilament Light (cat # M-1391-50, 2 µg per ~10<sup>6</sup> cells) for 30 minutes at room temperature, Secondary antibody: Goat anti-mouse PE labeled secondary antibody (1:100 fold dilution) with incubation for 20 minutes in dark at room temperature. Non-specific Control IgG, clone X63 (cat # M-1249-200) was used as negative control under same conditions (black dashed). Flow cytometry data and results were generated using Orflo Moxiflow™ instrument and protocols.

---

FOR RESEARCH USE ONLY