

## Mouse monoclonal antibody to Vimentin [VMT-24]: IgG

<b>Catalogue No.:</b>	M-1161-100
<b>Description:</b>	Vimentins are class-III intermediate filaments specific to mesenchymal tissue. Vimentin is an important cytoskeletal component responsible for maintaining cell integrity and has a probable role in the intracellular transport of proteins such as lipoproteins between the nucleus and plasma membrane. Immunohistochemical staining for Vimentin is characteristic of sarcomas.
<b>Batch No.:</b>	See product label
<b>Unit size:</b>	100 ug
<b>Antigen:</b>	Pig eye lens vimentin
<b>Clone:</b>	VMT-24
<b>Other Names:</b>	VIM;
<b>Accession:</b>	P08670 VIME_HUMAN; P02543 VIME_PIG;
<b>Produced in:</b>	Mouse
<b>Purity:</b>	IgG
<b>Applications:</b>	Immunohistochemistry (IHC) and Western Blotting (WB). A concentration of 0.5-1.0 ug/mL is recommended for WB. Human vimentin (precursor) has a predicted length of 466 residues and MW of 54 kDa. A concentration of 1.5-2.5 ug/mL is recommended to detect the protein in formalin fixed and paraffin embedded tissues. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
<b>Specificity:</b>	The specificity of this antibody has been confirmed by WB and IHC against the antigen.
<b>Cross-reactivity:</b>	Human; mouse; rat;
<b>Form:</b>	Lyophilized from 1.2% sodium acetate, 2mg BSA, 0.01mg NaN3
<b>Reconstitution:</b>	Reconstitute in 1 mL of PBS (pH 7.4) to achieve an antibody concentration of 100 ug/mL. Centrifuge to remove any insoluble material.
<b>Storage:</b>	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles
<b>Expiry Date:</b>	12 months after purchase

---

FOR RESEARCH USE ONLY