

Rabbit polyclonal antibody to Histone deacetylase 1 (1-14): Affinity purified

Catalogue No.:	R-1462-100
Description:	Histone deacetylase 1 is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression, and developmental events (Ref: SWISSPROT).
Batch No.:	See product label
Unit size:	100 ug
Antigen:	A synthetic peptide corresponding to a region (1-14 aa) from the N-terminus of human Histone deacetylase 1.
Antigen Location:	1-14 aa
Other Names:	HDAC1; HD1; RPD3L1;
Accession:	Q13547 HDAC1_HUMAN;
Produced in:	Rabbit
Applications:	Western Blotting (WB). A concentration of 1.0 ug/mL is recommended for WB. Human Histone deacetylase 1 has a predicted length of 482 amino acids and MW of 55 kDa. Biosensis recommends optimal dilutions/concentrations should be determined by the end user.
Specificity:	The specificity of this antibody has been confirmed by WB against the antigen.
Antibody Against:	Histone deacetylase 1
Cross-reactivity:	Human; rat; mouse;
Form:	Lyophilised with 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃
Appearance:	White powder
Reconstitution:	Reconstitute in 100 uL of sterile distilled water to achieve an antibody concentration of 1 mg/mL. Centrifuge to remove any insoluble material.
Storage:	At least 12 months after purchase at 2-8C (lyophilized formulations). After reconstitution, aliquot and store at -20C for a higher stability. Avoid freeze-thaw cycles.
Expiry Date:	12 months after purchase

FOR RESEARCH USE ONLY